

PATIENT GUIDE



FDA registered LidoPro[®] is formulated to meet the demands of minor to severe chronic pain sufferers. The patent-pending formulation delivers the highest concentration of safe and effective pain relieving ingredients available today, with several key stabilizing anti-irritant properties to uniformly deliver immediate local pain relief. LidoPro[®] is manufactured in accordance with FDA regulations by an FDA-approved manufacturer and is available under the supervision of licensed physicians, pharmacists, or other health care professionals.

PRODUCT INFORMATION

LidoPro[®] has been shown to reduce the body's inflammatory response, reduce pain, improve mobility, and increase musculoskeletal function. This special formulation utilizes the groundbreaking INUTEK[®] SP1 patented micro-emulsion technology to most effectively disperse and deliver the key active ingredient particles evenly, more rapidly, and deeper than alternative formulations. Quick and uniform delivery of the highly-concentrated ingredients allows for greater absorption and efficacy, while relief penetrates deeper into your muscles, joints, and nerves. LidoPro[®] contains many skin conditioners, such as Aloe Vera, Allantoin, and Chamomile for optimal comfort and is now available for hands-free use (see Directions for Use).

USES

Muscle Pain Due To: Myalgias, muscle strains or sprains, pulled muscles, muscle stiffness, muscle soreness, fibromyalgia, myofascial pain syndrome, exercise-induced myalgias, delayed onset myalgias, or piriformis syndrome.

Nerve Pain Due To: Neuropathies, neuralgia, neuritis, post herpetic neuralgia (shingles), occipital neuralgia, occipital neuritis, occipital neuropathy, peripheral neuralgia, peripheral neuritis, peripheral neuropathy, polyneuropathy, complex regional pain syndrome or CRPS, reflex sympathetic dystrophy or RSD, sudecks' atrophy, hand-shoulder syndrome, sciatica, meralgia paresthetica, intercostal neuralgia, intercostal neuritis, intercostal neuropathy, lumbar radiculopathy, lumbar radiculitis, or pinched nerves.

Joint Pain Due To: Osteoarthritis (OA), rheumatoid arthritis (RA), juvenile rheumatoid arthritis (JRA), degenerative joint disease (DJD), psoriatic arthritis, gouty arthritis, cervical facet joint disease, lumbar facet joint disease, bursitis, carpal tunnel syndrome, arthritis, or impingement syndrome of the shoulder.

Ligament Pain Due To: Sprains or strains, tendonitis, tendonopathy, epichondylitis, or costochondritis.

Spinal, Neck or Back Pain Due To: Bulging disc, herniated disc, slipped disc, degenerative disc disease, degenerative joint disease, facet joint arthritis, musculoligamentous strain, sacroiliitis, scoliosis, spinal stenosis, spondylolithesis, or spondylosis.

Body Pains, including: Arm, shoulder, elbow, wrist, hand, chest, foot, ankle, toe, head, hip, knee, leg, back, or neck.

DIRECTIONS FOR USE

Use LidoPro[®] only as directed by healthcare professional. Shake well before each use. Before first use, unscrew the Applicator unit and remove the safety seal from the bottom of the tube. Replace Applicator unit so it is secure. Pull off Applicator cap. With Applicator facing upward, gently squeeze tube until approximately a dime-sized quantity of ointment emerges on Applicator surface. Use tube and Applicator to delicately rub ointment into affected area. Do not rub vigorously or use Applicator to massage ointment into skin. Use as needed for effective relief, but no more than four applications per day. LidoPro[®] can still be applied by hand. Simply unscrew the Applicator unit and administer by hand. Make sure to wash hands after using them to apply ointment. Washing hands after every use — including use of Applicator — is recommended. When using this product, avoid contact with eyes, genitals, and other mucous membranes. If eye contact occurs, rinse thoroughly with cold water and call a doctor. If swallowed, call poison control immediately. For best results, continue use for one to four weeks. Stop use if itching or rash occurs. Keep away from children. If pregnant or nursing, please consult your doctor or physician. Avoid storing product in direct sunlight. Protect product from excessive moisture. If pregnant or nursing, please consult your doctor or physician. Avoid storing product in direct sunlight. Protect product from excessive moisture. Store at 68-77°F (20-25°C).

Do Not Use • on open wounds, cuts, damaged or infected skin • with bandage or a heating pad • if condition worsens or symptoms persist for more than 7 days • if you are allergic to any ingredients, PABA, aspirin products or sulfa • excessive skin irritation occurs • in large amounts or over raw or blistered skin • if safety seal has been tampered • while in hot • tub or sauna • if you are taking Warfarin or Coumadin